

ON TORIC LOCALLY CONFORMALLY KÄHLER MANIFOLDS

FARID MADANI, ANDREI MOROIANU, MIHAELA PILCA

ABSTRACT. We study compact toric strict locally conformally Kähler manifolds. We show that the Kodaira dimension of the underlying complex manifold is $-\infty$ and that the only compact complex surfaces admitting toric strict locally conformally Kähler metrics are the diagonal Hopf surfaces. We also show that every toric Vaisman manifold has lcK rank 1 and is isomorphic to the mapping torus of an automorphism of a toric compact Sasakian manifold.

1. INTRODUCTION

When searching for the *best* Hermitian metrics on a compact complex manifold (M, J) , one is naturally led to consider Kähler metrics. However, there are well-known topological obstructions which severely restrict the class of compact complex manifolds carrying such metrics. A more general class of compatible metrics, which was introduced by I. Vaisman [21] in the 70's, are the *locally conformally Kähler (lcK)* metrics. These are characterized by the condition that around any point in M , the metric g can be conformally rescaled to a Kähler metric. If this metric can be globally defined, the structure is globally conformally Kähler (gcK), otherwise it is called *strict* lcK. The topological obstructions imposed by the existence of lcK metrics are less restrictive than in the Kähler case. In complex dimension 2, for instance, it was widely believed that any compact complex surface admits an lcK metric, until F. Belgun [1] proved that some Inoue surfaces do not admit any lcK metric.

In this paper we investigate compact strict lcK manifolds admitting an effective torus action of maximal dimension by twisted Hamiltonian biholomorphisms. Such structures are called toric lcK and were introduced in [17].

The paper is organized as follows: after some preliminaries on lcK manifolds, we study the properties of their automorphism group in Section 3 and give several general results about lifts of group actions on lcK manifolds in Section 4.

In Section 5 we use the special decomposition of the space of harmonic 1-forms on compact Vaisman manifolds to show that every toric Vaisman manifold has first Betti number $b_1 = 1$ and is thus isomorphic to the mapping torus of an automorphism of a toric compact Sasakian manifold.

2010 *Mathematics Subject Classification.* 53A30, 53B35, 53C25, 53C29, 53C55.

Key words and phrases. locally conformally Kähler structure, Hopf surface, toric manifold, twisted Hamiltonian.

In Section 6 we show that the Kodaira dimension of every compact complex manifold admitting a toric strict lcK structure is $-\infty$ (see Theorem 6.1).

Using this result and the Kodaira classification of non-Kählerian compact complex surfaces, we show in Theorem 7.2 that the only compact complex surfaces with a toric strict lcK metric are the *diagonal Hopf surfaces*. Indeed, lcK metrics on these manifolds have been constructed by P. Gauduchon and L. Ornea, [8], and they turn out to admit toric T^2 -actions.

Acknowledgments. This work was supported by the Procope Project No. 32977YJ and by the SFB 1085. We thank Nicolina Istrati for pointing out to us an error in a preliminary version of the paper and for useful suggestions.

2. PRELIMINARIES ON LcK MANIFOLDS

Let (M^{2n}, J) be a (connected) complex manifold of complex dimension $n \geq 2$. A Kähler structure on (M, J) is a Riemannian metric g compatible with J (in the sense that J is skew-symmetric with respect to g), and such that the 2-form $\Omega := g(J\cdot, \cdot)$ is closed.

Two Kähler structures g and g' on (M, J) in the same conformal class are necessarily homothetic. Indeed, if $g' = fg$ for some positive function f , the associated 2-forms are related by $\Omega' = f\Omega$, thus $0 = d\Omega' = df \wedge \Omega + f d\Omega = df \wedge \Omega$, which shows that $df = 0$, since the wedge product with the non-degenerate form Ω is injective on 1-forms.

A locally conformally Kähler (lcK) structure on (M, J) is a Riemannian metric g which is locally conformally Kähler in the sense that there exists an open covering $\{U_\alpha\}_{\alpha \in \mathcal{A}}$ of M and smooth maps $\varphi_\alpha \in C^\infty(U_\alpha)$ such that $(U_\alpha, J, e^{-\varphi_\alpha}g)$ is Kähler. This definition is clearly independent on the metric g in its conformal class $[g]$, so one usually refers to $(J, [g])$ as being an lcK structure.

Since $e^{-\varphi_\alpha}g$ and $e^{-\varphi_\beta}g$ are Kähler metrics in the same conformal class on $U_\alpha \cap U_\beta$, we deduce from the above remark that $\varphi_\alpha - \varphi_\beta$ is locally constant, and thus $d\varphi_\alpha = d\varphi_\beta$ on $U_\alpha \cap U_\beta$. This shows the existence of a closed 1-form θ , called the *Lee form* of the lcK structure (J, g) , such that $\theta|_{U_\alpha} = d\varphi_\alpha$. If $\Omega := g(J\cdot, \cdot)$ denotes like before the 2-form associated to g and J , then by definition $e^{-\varphi_\alpha}\Omega$ is a closed form on U_α , thus

$$0 = d(e^{-\varphi_\alpha}\Omega) = e^{-\varphi_\alpha}(-d\varphi_\alpha \wedge \Omega + d\Omega) = e^{-\varphi_\alpha}(-\theta \wedge \Omega + d\Omega),$$

showing that $d\Omega = \theta \wedge \Omega$ everywhere on M . If the lcK metric is changed by a conformal factor, $g' := e^{-\varphi}g$, the corresponding Lee form satisfies $\theta' = \theta - d\varphi$.

If the Lee form vanishes, g is Kähler, if it is exact, $\theta = d\varphi$, then g is globally conformal to the Kähler metric $g' = e^{-\varphi}g$, and if it is parallel, g is called *Vaisman*. In this paper we will always assume that the lcK structure is *strict*, in the sense that its Lee form is not exact. This definition clearly does not depend on the choice of the metric g in its conformal class.

Let now \widetilde{M} be the universal covering of an lcK manifold (M, J, g, θ) , endowed with the pull-back lcK structure $(\widetilde{J}, \widetilde{g}, \widetilde{\theta})$. Since \widetilde{M} is simply connected, $\widetilde{\theta}$ is exact, *i.e.* $\widetilde{\theta} = d\varphi$, and by the above considerations, the metric $g^K := e^{-\varphi}\widetilde{g}$ is Kähler. The group $\pi_1(M)$ acts on $(\widetilde{M}, \widetilde{J}, g^K)$ by holomorphic homotheties. Indeed, since $\widetilde{\theta}$ is $\pi_1(M)$ -invariant, there exists a

group morphism $\rho: \pi_1(M) \rightarrow (\mathbb{R}, +)$, $\gamma \mapsto c_\gamma$, such that

$$(1) \quad \gamma^* \varphi = \varphi + c_\gamma, \quad \forall \gamma \in \pi_1(M).$$

We thus have $\gamma^* g^K = e^{-c_\gamma} g^K$, for every $\gamma \in \pi_1(M)$. The Kähler structure (\tilde{J}, g^K) is tautologically invariant by $\ker \rho$ and defines a Kähler structure denoted (\hat{J}, g^K) on $\widehat{M} := \widetilde{M} / \ker \rho$. The Kähler manifold $(\widehat{M}, \hat{J}, g^K)$ is called the *minimal covering* of (M, J, g) (and it actually only depends on the conformal class $[g]$).

We thus obtain the following sequence of Galois coverings

$$\widetilde{M} \xrightarrow{\ker \rho} \widehat{M} \xrightarrow{\Gamma} M,$$

where $\Gamma := \pi_1(M) / \ker \rho$. Since ρ induces an injective group homomorphism $\Gamma \rightarrow \mathbb{R}_+^*$, it follows that the automorphism group Γ of the minimal covering is isomorphic to a free Abelian group \mathbb{Z}^k , for some $k \in \mathbb{N}$. The integer k is called the *rank* of the lcK structure.

A useful tool in lcK geometry is the so-called *Gauduchon metric*. If $(M, J, [g])$ is a compact lcK manifold, the Gauduchon metric $g_0 \in [g]$ is characterized by the fact that its Lee form is co-closed: $\delta^{g_0} \theta_0 = 0$, and it is unique, up to a positive factor, in the given conformal class, [7].

The so-called *twisted differential* of an lcK structure (M, J, g, θ) is defined as follows: $d^\theta: \Omega^*(M) \rightarrow \Omega^{*+1}(M)$, $d^\theta \alpha := d\alpha - \theta \wedge \alpha$. Note that $d\theta = 0$ implies $d^\theta \circ d^\theta = 0$. However, d^θ does not satisfy the Leibniz rule. If $\alpha \in \Omega^k(M)$ and $\beta \in \Omega^\ell(M)$, then the following relation holds:

$$d^\theta(\alpha \wedge \beta) = d^\theta \alpha \wedge \beta + (-1)^k \alpha \wedge d^\theta \beta + \theta \wedge \alpha \wedge \beta.$$

In particular, for $f \in \mathcal{C}^\infty(M)$, we have

$$(2) \quad d^\theta(f\beta) = df \wedge \beta + f d^\theta \beta.$$

Lemma 2.1. *On a strict lcK manifold $(M, J, [g])$, the twisted differential acting on functions is injective, i.e. $\ker(d^\theta: \mathcal{C}^\infty(M) \rightarrow \Omega^1(M)) = \{0\}$.*

Proof. Let $f \in \ker(d^\theta)$, so f satisfies the linear first order differential equation $df - f\theta = 0$. If f does not vanish at any point of the manifold, then $\theta = d \log |f|$, which is not possible, since the lcK manifold is assumed to be strict. Thus, f vanishes at some point, which implies that $f \equiv 0$, by uniqueness of the solution. \square

3. AUTOMORPHISMS OF LCK MANIFOLDS

An automorphism of an lcK manifold $(M, J, [g])$ is a conformal biholomorphism. We denote by $\text{Aut}(M, J, [g])$ the group of all automorphisms and by $\mathfrak{aut}(M, J, [g])$ its Lie algebra. In this section we establish a few properties of the automorphisms of lcK manifolds, that will be used in the sequel. Let us recall the following notions:

Definition 3.1. Let (M, J, g, θ) be an lcK manifold. A vector field X on M is called *twisted Hamiltonian* if there exists $h_X \in \mathcal{C}^\infty(M)$, such that $X \lrcorner \Omega = d^\theta h_X$. The space of all such vector fields on M is denoted by $\mathfrak{ham}^\theta(M)$. An action of a Lie group on M is called *twisted Hamiltonian* if all its fundamental vector fields belong to $\mathfrak{ham}^\theta(M)$.

Definition 3.2. An lcK manifold $(M^{2n}, J, [g])$ equipped with an effective holomorphic and twisted Hamiltonian action of the standard (real) n -dimensional torus T^n , is called *toric* lcK.

Next we give a class of examples of toric Vaisman manifolds. For this purpose, let us recall the definition of a Sasakian manifold. A *Sasakian structure* on a Riemannian manifold (S, g_S) is a complex structure \tilde{J} on $\mathbb{R} \times S$, such that the cone metric $g^K := e^{-2t}(dt^2 + g_S)$ is Kähler with respect to \tilde{J} , and for each $\lambda \in \mathbb{R}$, the homothety $(t, w) \mapsto (t + \lambda, w)$ is holomorphic.

Example 3.3. An example of a toric lcK manifold is the Hopf manifold $S^1 \times S^{2n-1}$, whose natural Vaisman structure is toric, as noticed in [17, Example 4.8]. More generally, a class of toric Vaisman manifolds can be constructed as follows. Given a compact toric Hodge manifold N of complex dimension $n - 1$, we consider the total space of the S^1 -bundle S corresponding to the integral cohomology class of its Kähler form, which carries a Sasakian metric g_S . Hence the product $\mathbb{R} \times S$ is endowed with a Kähler structure $(g^K := e^{-2t}(dt^2 + g_S), \tilde{J})$. The action of \mathbb{Z} by translations on \mathbb{R} and extended trivially on S is holomorphic and isometric with respect to the gcK structure $(\tilde{J}, \tilde{g} := e^{2t}g^K, 2dt)$, which thus projects onto a Vaisman structure (J, g, θ) on the compact manifold $S^1 \times S$. By construction, N is the quotient $(S^1 \times S)/\{\theta^\sharp, J\theta^\sharp\}$ and its toric Kähler structure is the projection of (J, g) . According to [17, Theorem 5.1], this Vaisman structure on $S^1 \times S$ is toric.

We continue by proving some preliminary results.

Lemma 3.4. *Let (M, J, g, θ) be an lcK manifold with fundamental 2-form Ω . There exists a morphism of Lie algebras $\sigma: \mathbf{aut}(M, J, [g]) \rightarrow \mathbb{R}$, such that for any $X \in \mathbf{aut}(M, J, [g])$, we have*

$$(3) \quad d^\theta(X \lrcorner \Omega) = \sigma(X)\Omega,$$

$$(4) \quad \mathcal{L}_X g = (\theta(X) + \sigma(X))g.$$

Proof. For $X \in \mathbf{aut}(M, J, [g])$, there exists $f_X \in C^\infty(M)$, such that $\mathcal{L}_X \Omega = f_X \Omega$. By the Cartan formula, we have

$$(5) \quad (f_X - \theta(X))\Omega = \mathcal{L}_X \Omega - \theta(X)\Omega = d(X \lrcorner \Omega) - \theta \wedge (X \lrcorner \Omega) = d^\theta(X \lrcorner \Omega).$$

Since $(d^\theta)^2 = 0$ and Ω is d^θ -closed, it follows by (2) that $d(f_X - \theta(X)) \wedge \Omega = 0$, which by the injectivity of $\Omega \wedge \cdot$ on 1-forms implies that $d(f_X - \theta(X)) = 0$. Hence there exists $\sigma(X) \in \mathbb{R}$, such that $f_X = \theta(X) + \sigma(X)$. Since X is a holomorphic vector field, it follows that $\mathcal{L}_X g = (\theta(X) + \sigma(X))g$.

It remains to show that σ is a Lie algebra morphism, *i.e.* that $\sigma([X, Y]) = 0$, for all $X, Y \in \mathbf{aut}(M, J, [g])$. Since θ is closed and $\sigma(X)$ and $\sigma(Y)$ are constant functions, we obtain

$$(6) \quad \theta([X, Y]) = X(\theta(Y)) - Y(\theta(X)) = X(f_Y - \sigma(Y)) - Y(f_X - \sigma(X)) = X(f_Y) - Y(f_X).$$

On the other hand, we compute

$$\mathcal{L}_{[X, Y]} \Omega = \mathcal{L}_X \mathcal{L}_Y \Omega - \mathcal{L}_Y \mathcal{L}_X \Omega = \mathcal{L}_X (f_Y \Omega) - \mathcal{L}_Y (f_X \Omega) = (X(f_Y) - Y(f_X))\Omega,$$

so that $f_{[X, Y]} = X(f_Y) - Y(f_X)$. By (6), it follows that $\sigma([X, Y]) = f_{[X, Y]} - \theta([X, Y]) = 0$. \square

Remark 3.5. Let $(M, J, [g])$ be an lcK manifold. On its universal covering \widetilde{M} endowed with the Kähler structure (g^K, Ω^K) , Lemma 3.4 reads

$$(7) \quad d(\widetilde{X} \lrcorner \Omega^K) = \sigma(X)\Omega^K,$$

$$(8) \quad \mathcal{L}_{\widetilde{X}}g^K = \sigma(X)g^K,$$

for each vector field X on M , whose lift on \widetilde{M} is denoted by \widetilde{X} .

We denote by $\text{Aut}_s(M, J, [g])$ the subgroup of $\text{Aut}(M, J, [g])$ consisting of automorphisms whose lifts to \widetilde{M} are isometries with respect to the Kähler metric g^K . By (8), its Lie algebra is $\mathfrak{aut}_s(M, J, [g]) := \ker \sigma$. The elements of $\text{Aut}_s(M, J, [g])$ are called *special* automorphisms of the lcK structure $(M, J, [g])$.

Lemma 3.6. *On an lcK manifold $(M^{2n}, J, [g])$, the following inclusion holds:*

$$(9) \quad \mathfrak{ham}^\theta(M) \cap \mathfrak{hol}(M) \subseteq \mathfrak{aut}_s(M, J, [g]),$$

where $\mathfrak{hol}(M)$ denotes the set of holomorphic vector fields on (M, J) .

Proof. Let $X \in \mathfrak{ham}^\theta(M) \cap \mathfrak{hol}(M)$. Then there exists $h_X \in \mathcal{C}^\infty(M)$, such that $X \lrcorner \Omega = d^\theta h_X$. We now compute using Cartan's formula and the fact that $d = d^\theta + \theta \wedge$.

$$(10) \quad \begin{aligned} \mathcal{L}_X \Omega &= d^\theta(X \lrcorner \Omega) + \theta \wedge (X \lrcorner \Omega) + X \lrcorner d^\theta \Omega + X \lrcorner (\theta \wedge \Omega) \\ &= \theta \wedge (X \lrcorner \Omega) + X \lrcorner (\theta \wedge \Omega) = \theta(X)\Omega. \end{aligned}$$

Since X is a holomorphic vector field, it follows that $\mathcal{L}_X g = \theta(X)g$. Hence, $X \in \mathfrak{aut}(M, J, [g])$. Moreover, (4) implies that $X \in \ker \sigma$. \square

It turns out that this inclusion is even an equality under some additional assumption:

Lemma 3.7. *Let $(M^{2n}, J, [g])$ be a compact lcK manifold. Assume that the minimal covering of M coincides with its universal covering. Then $\mathfrak{ham}^\theta(M) \cap \mathfrak{hol}(M) = \mathfrak{aut}_s(M, J, [g])$.*

Proof. Let $\Omega^K = e^{-\varphi} \pi^* \Omega$ be the fundamental form of the Kähler structure on the universal covering of M , where $\pi^* \theta = d\varphi$. Let us fix some $X \in \mathfrak{aut}_s(M, J, [g])$. By (7), the following equality holds $d(\widetilde{X} \lrcorner \Omega^K) = 0$, where \widetilde{X} is the lift of X to \widetilde{M} . Thus, there exists a function $h \in \mathcal{C}^\infty(\widetilde{M})$, such that $dh = \widetilde{X} \lrcorner \Omega^K$.

The hypothesis implies that every non-trivial element in $\pi_1(M)$ acts by a strict homothety on (\widetilde{M}, g^K) . Thus $\pi_1(M)$ is commutative, since a commutator of homotheties is an isometry. For each $\gamma \in \pi_1(M)$, we have $\gamma^* \varphi = \varphi + c_\gamma$, where $\gamma \mapsto c_\gamma$ is the group morphism defined in (1). Hence, we have $\gamma^* \Omega^K = e^{-c_\gamma} \Omega^K$, which implies $\gamma^* dh = e^{-c_\gamma} dh$, or equivalently $d(\gamma^* h - e^{-c_\gamma} h) = 0$. Therefore, there exists a constant λ_γ , such that $\gamma^* h - e^{-c_\gamma} h = \lambda_\gamma$. Let us fix some $\gamma_0 \in \pi_1(M) \setminus \{\text{id}\}$. By replacing h with $h - \frac{\lambda_{\gamma_0}}{1 - e^{-c_{\gamma_0}}}$, we may assume $\lambda_{\gamma_0} = 0$. We now prove that in fact $\lambda_\gamma = 0$, for all $\gamma \in \pi_1(M)$. Comparing the following two relations

$$\begin{aligned} \gamma^* \gamma_0^* h &= \gamma^*(e^{-c_{\gamma_0}} h) = e^{-c_{\gamma_0}} (e^{-c_\gamma} h + \lambda_\gamma), \\ \gamma_0^* \gamma^* h &= \gamma_0^*(e^{-c_\gamma} h + \lambda_\gamma) = e^{-c_{\gamma_0} - c_\gamma} h + \lambda_\gamma, \end{aligned}$$

and using the commutativity of $\pi_1(M)$, we obtain $\lambda_\gamma(1 - e^{-c_\gamma}) = 0$. As $c_\gamma \neq 0$, it follows that $\lambda_\gamma = 0$, for any $\gamma \in \pi_1(M)$. Concluding, we have shown that

$$\gamma^*h = e^{-c_\gamma}h, \quad \forall \gamma \in \pi_1(M).$$

It follows that the function $e^\varphi h$ projects onto M . Moreover, on \widetilde{M} we have

$$d^\theta(e^\varphi h) = e^\varphi(h\theta + dh) - e^\varphi h\theta = e^\varphi \widetilde{X} \lrcorner \Omega^K = \widetilde{X} \lrcorner \pi^* \Omega.$$

Thus, X is a twisted Hamiltonian vector field on M , whose Hamiltonian function is the projection of $e^\varphi h$ onto M . \square

Proposition 3.8. *Let $(M^{2n}, J, [g])$ be a compact lcK manifold. The following assertions hold:*

- (i) *Every vector field in $\mathbf{aut}(M, J, [g])$ is a Killing vector field for the Gauduchon metric $g_0 \in [g]$.*
- (ii) *If θ_0 denotes the Lee form of the Gauduchon metric, we have $\mathbf{aut}_s(M, J, [g]) \subseteq \ker \theta_0$.*
- (iii) *For any $X \in \mathbf{aut}_s(M, J, [g])$, its lift \widetilde{X} , resp. \widehat{X} , to the universal (resp. minimal) covering of $(M, J, [g])$ is a Killing vector field for the Kähler metric g^K .*

Proof. (i) It suffices to prove that any $\psi \in \text{Aut}(M^{2n}, J, [g])$ is an isometry of g_0 . We first notice that the metric $\psi^*g_0 \in [\psi^*g]$ is also a Gauduchon metric. By definition, ψ preserves the conformal class, so $[\psi^*g] = [g]$. Since in any conformal class, a Gauduchon metric is unique up to a positive constant, there exists $c > 0$, such that $\psi^*g_0 = cg_0$. This means that ψ is a homothety. However, on a compact manifold any homothety is already an isometry. Therefore, ψ is an isometry for g_0 .

(ii) Follows from (i) and (4).

(iii) Follows from (8). \square

Proposition 3.9. *Let $(M^{2n}, J, [g])$ be a compact strict lcK manifold endowed with an effective twisted Hamiltonian holomorphic T^m -action. Then the orbits of the action are isotropic with respect to the fundamental 2-form Ω and $m \leq n$.*

Proof. Let X, Y be two fundamental vector fields associated to the given T^m -action on M . Lemma 3.6 implies that $X, Y \in \mathbf{aut}_s(M)$. We then compute using Lemma 3.4 and the fact that $[X, Y] = 0$:

$$\begin{aligned} d(\Omega(X, Y)) &= d(Y \lrcorner \Omega(X)) = \mathcal{L}_Y(\Omega(X)) - Y \lrcorner d(\Omega(X)) = (\mathcal{L}_Y \Omega)(X) - Y \lrcorner (\mathcal{L}_X \Omega - X \lrcorner d\Omega) \\ &= \theta(Y)\Omega(X) - \theta(X)\Omega(Y) + Y \lrcorner X \lrcorner (\theta \wedge \Omega) = \Omega(X, Y)\theta. \end{aligned}$$

This shows that $d^\theta(\Omega(X, Y)) = 0$. By Lemma 2.1, we conclude that $\Omega(X, Y) = 0$, so each T^m -orbit is isotropic with respect to Ω and hence has dimension at most n . On the other hand, a known consequence of the principal orbit theorem (see *e.g.* [6, Theorem 2.8.5]) is that any T^m acting effectively on a compact manifold, acts freely on its principal orbit. Thus, it follows that $m \leq n$. \square

4. GROUP ACTIONS ON LCK MANIFOLDS

We start with a few general lemmas about lifts of group actions from the basis to the total space of a covering. Let G be a connected Lie group acting smoothly on a connected manifold M , and let $\pi : \widehat{M} \rightarrow M$ be a covering with connected covering space. For every $x \in M$ we denote by $f_x : G \rightarrow M$ the map $f_x(a) := a \cdot x$.

Lemma 4.1. *The action of G on M lifts to an action of G on \widehat{M} if and only if for each $x \in M$ and $\hat{x} \in \widehat{M}$ with $\pi(\hat{x}) = x$, the map f_x lifts to a smooth map $f_{\hat{x}} : G \rightarrow \widehat{M}$ which maps 1_G to \hat{x} .*

Proof. One direction is clear. Conversely, we define the map $f : G \times \widehat{M} \rightarrow \widehat{M}$ by $(a, \hat{x}) \mapsto a \cdot \hat{x} := f_{\hat{x}}(a)$. Since $\pi \circ f(a, \hat{x}) = a \cdot \pi(\hat{x})$, the map f is smooth. Moreover, for every $a, b \in G$ and $\hat{x} \in \widehat{M}$ we have

$$\pi(a \cdot (b \cdot \hat{x})) = a \cdot \pi(b \cdot \hat{x}) = a \cdot (b \cdot \pi(\hat{x})) = (ab) \cdot \pi(\hat{x}) = \pi((ab) \cdot \hat{x}),$$

thus showing that $a \cdot (b \cdot \hat{x})$ and $(ab) \cdot \hat{x}$ belong to the same fibre. By continuity of the map f , the set of points (a, b, \hat{x}) such that $a \cdot (b \cdot \hat{x}) = (ab) \cdot \hat{x}$ is open, closed and contains $1_G \times G \times \widehat{M} \subset G \times G \times \widehat{M}$. Since $G \times G \times \widehat{M}$ is connected, the above relation holds identically, thus showing that f is a group action. \square

Corollary 4.2. *If $G = G_1 \times G_2$ acts on M and the action of G_i on M lifts to an action of G_i on \widehat{M} for $i = 1, 2$, then the action of G on M lifts to an action of G on \widehat{M} .*

Proof. We apply the previous lemma, noting that for each $x \in M$ and $\hat{x} \in \widehat{M}$ the map f_x lifts to the map $f_{\hat{x}}(a_1, a_2) := a_1 \cdot (a_2 \cdot \hat{x})$. \square

Lemma 4.3. *Assume in addition that \widehat{M} is a Galois covering of M whose automorphism group has no torsion and let $p : \widehat{G} \rightarrow G$ be a finite covering. If the induced action of \widehat{G} on M lifts to \widehat{M} , then the action of G lifts to \widehat{M} .*

Proof. We denote by K the image of $\pi_1(\widehat{M})$ in $\pi_1(M)$. The hypothesis shows that K is a normal subgroup of $\pi_1(M)$, and $\pi_1(M)/K$ has no element of finite order (except the identity).

For every $x \in M$ we consider as before the map $f_x : G \rightarrow M$, $a \mapsto a \cdot x$ and denote by $\hat{f}_x := f_x \circ p : \widehat{G} \rightarrow M$. By the classical covering lemma, f_x lifts to \widehat{M} if and only if $(f_x)_*(\pi_1(G)) \subset K$. By assumption, \hat{f}_x lifts to \widehat{M} , so we have $(\hat{f}_x)_*(\pi_1(\widehat{G})) \subset K$, whence $(f_x)_*(p_*(\pi_1(\widehat{G}))) \subset K$. The map f_* thus induces a group morphism

$$\rho : \pi_1(G)/p_*(\pi_1(\widehat{G})) \rightarrow \pi_1(M)/K.$$

On the other hand, $p_*(\pi_1(\widehat{G}))$ has finite index in $\pi_1(G)$ since $\widehat{G} \rightarrow G$ is a finite covering, and $\pi_1(M)/K$ has no torsion, so the morphism ρ vanishes identically. This shows that $(f_x)_*(\pi_1(G)) \subset K$, so f_x lifts to \widehat{M} . We conclude by Lemma 4.1. \square

Proposition 4.4. *Let (M^{2n}, J) be a compact complex manifold carrying an lcK structure (J, g, θ) with minimal covering \widehat{M} . Let G be a compact connected Lie group acting holomorphically on M . Then the following assertions hold:*

- (i) *If the action of G lifts to \widehat{M} , then there exists an lcK structure (J, g', θ') on M , such that $G \subseteq \text{Aut}_s(M, J, [g'])$.*
- (ii) *If $G \subseteq \text{Aut}_s(M, J, [g])$, then the action of G lifts to \widehat{M} .*

Proof. (i) Let Γ denote the automorphism group of the covering $\pi : \widehat{M} \rightarrow M$. The pull-back $\pi^*\theta$ of the Lee form to \widehat{M} is exact, so there exists a function φ on \widehat{M} such that $\pi^*\theta = d\varphi$. Since $\pi^*\theta$ is Γ -invariant, there exists a group morphism $\Gamma \rightarrow (\mathbb{R}, +)$, $\gamma \mapsto c_\gamma$, such that $\gamma^*\varphi = \varphi + c_\gamma$ for every $\gamma \in \Gamma$, similar to (1). The Kähler metric $g^K := e^{-\varphi}\pi^*g$ is then Γ -equivariant, in the sense that $\gamma^*g^K = e^{-c_\gamma}g^K$, for every $\gamma \in \Gamma$.

We claim that G and Γ commute. By assumption, for every $a \in G$ and $\hat{x} \in \widehat{M}$, we have $\pi(a\hat{x}) = a\pi(\hat{x})$. Consequently, for every $\gamma \in \Gamma$ we obtain

$$\pi(a\gamma\hat{x}) = a\pi(\gamma\hat{x}) = a\pi(\hat{x}) = \pi(a\hat{x}).$$

Since G is connected and Γ is discrete, this shows that for every $\gamma \in \Gamma$ there exists $\gamma' \in \Gamma$ such that $a\gamma = \gamma'a$ for every $a \in G$. Taking $a = \text{id}$ shows that $\gamma = \gamma'$, thus proving our claim.

Let now $d\mu$ denote the Haar measure of G (normalized such that G has unit volume) and let $\widehat{\varphi}$ and $\widehat{g^K}$ be the average on \widehat{M} over G of φ and g^K respectively:

$$\widehat{\varphi} = \int_G a^*\varphi d\mu(a), \quad \widehat{g^K} = \int_G a^*g^K d\mu(a).$$

Clearly $\widehat{g^K}$ is still Kähler, and by construction, taking into account the commutation of G and Γ , we see that the function $\widehat{\varphi}$ and the metric $\widehat{g^K}$ are G -invariant and Γ -equivariant:

$$\gamma^*\widehat{\varphi} = \widehat{\varphi} + c_\gamma, \quad \gamma^*\widehat{g^K} = e^{-c_\gamma}\widehat{g^K}, \quad \forall \gamma \in \Gamma.$$

It follows that the metric $e^{\widehat{\varphi}}\widehat{g^K}$ projects onto a G -invariant lcK metric g' on M , whose corresponding Lee form θ' satisfies $\theta'(X) = 0$, for any fundamental vector field X of the G -action. Since such an X is also a Killing vector field with respect to g' , we obtain by (4) that $G \subseteq \text{Aut}_s(M, J, [g'])$.

(ii) Any compact Lie group is a finite quotient of $T^k \times G_s$, for some $k \in \mathbb{N}$ and G_s a simply connected compact Lie group. By Lemma 4.1, the action of G_s lifts to \widehat{M} , so using Corollary 4.2 and Lemma 4.3, it suffices to show that if $T^k \subseteq \text{Aut}_s(M, J, [g])$ on a compact lcK manifold $(M^{2n}, J, [g])$, then its action lifts to the minimal covering \widehat{M} . By Corollary 4.2 again, we may assume $k = 1$. Equivalently, we need to show that the lift of every $X \in \mathfrak{aut}_s(M, J, [g])$ with closed orbits on M has closed orbits on \widehat{M} .

By Proposition 3.8, X is Killing with respect to the Gauduchon metric g_0 on M , and its lift \widehat{X} to \widehat{M} is Killing with respect to both π^*g_0 and g^K . Since the flow of X is equal to the identity at some time $t_0 > 0$, it follows that the flow of \widehat{X} at time t_0 is an automorphism

of the covering \widehat{M} , and at the same time an isometry of g^K , as \widehat{X} preserves g^K . On the other hand, by definition of the minimal covering, the only automorphism of \widehat{M} which is an isometry of g^K is the identity, thus showing that \widehat{X} has closed orbits. \square

A straightforward consequence of Proposition 4.4 is the following:

Corollary 4.5. *Let (M^{2n}, J) be a compact complex manifold and let G be a compact Lie group acting holomorphically on M . If the G -action does not lift to any of the non-compact coverings of M , then there exists no lcK structure $(J, [g])$ on M , such that $G \subseteq \text{Aut}_s(M, J, [g])$.*

5. TORIC VAISMAN MANIFOLDS

In this section, we investigate toric compact Vaisman manifolds and show that each such manifold is obtained as the mapping torus of a compact toric Sasakian manifold.

Without loss of generality, we assume that the norm of the Lee form of a Vaisman structure equals 1. Recall that on a Vaisman manifold the Lee and anti-Lee vector fields θ^\sharp and $J\theta^\sharp$ are Killing and holomorphic. We will need in the sequel the following elementary observation:

Lemma 5.1. *For any Killing vector field X and any harmonic 1-form α on a compact connected Riemannian manifold, the following identity holds $\mathcal{L}_X\alpha = d(\alpha(X)) = 0$. In particular, $\alpha(X)$ is constant.*

Proof. Since the manifold is compact, α is closed and co-closed. Cartan's formula then yields $\mathcal{L}_X\alpha = d(\alpha(X))$. On the other hand, the 1-form $\mathcal{L}_X\alpha$ is harmonic, as X is a Killing vector field. By the Hodge decomposition theorem on a compact Riemannian manifold, the only harmonic and exact differential form is the trivial one. Thus, $\mathcal{L}_X\alpha = d(\alpha(X)) = 0$. \square

The following result is folklore, see *e.g.* [11], [19], [20], [23]. For the convenience of the reader we give here an elementary proof.

Lemma 5.2. *Let (M^{2n}, J, g, θ) be a compact Vaisman manifold of complex dimension $n \geq 2$.*

- (i) *If $\alpha \in \Omega^1(M)$ is a harmonic form pointwise orthogonal on θ , then $J\alpha$ is also harmonic and pointwise orthogonal on θ .*
- (ii) *The following decomposition of the space of harmonic 1-forms holds:*

$$(11) \quad \mathcal{H}^1(M) = \text{span}\{\theta\} \oplus \mathcal{H}_0^1(M),$$

where each element of $\mathcal{H}_0^1(M)$ is pointwise orthogonal on θ and $\mathcal{H}_0^1(M)$ is J -invariant.

Proof. (i) Lemma 5.1 applied to the harmonic form α and the Killing vector fields θ^\sharp and $J\theta^\sharp$ yields

$$(12) \quad \mathcal{L}_{\theta^\sharp}\alpha = 0, \quad \mathcal{L}_{J\theta^\sharp}\alpha = d(\alpha(J\theta^\sharp)) = 0$$

and the function $\langle J\theta, \alpha \rangle =: c$ is constant on M . Furthermore, by Cartan's formula and using the fact that θ^\sharp is a holomorphic vector field, we obtain

$$(13) \quad \theta^\sharp \lrcorner dJ\alpha = d\langle J\alpha, \theta \rangle + \theta^\sharp \lrcorner dJ\alpha = \mathcal{L}_{\theta^\sharp}(J\alpha) = 0.$$

Recall that on the universal covering \widetilde{M} , endowed with the pull-back lcK structure $(\widetilde{J}, \widetilde{g}, \widetilde{\theta})$, the metric \widetilde{g} and the Kähler metric g^K are conformally related by $g^K = e^{-\varphi}\widetilde{g}$, with $d\varphi = \widetilde{\theta}$. Hence, their corresponding Laplace operators acting on a 1-form β on \widetilde{M} are related as follows (see for instance [3, Theorem 1.159]):

$$(14) \quad \Delta^K \beta = e^\varphi \left(\widetilde{\Delta} \beta + (n-1)d(\langle \beta, \widetilde{\theta} \rangle) + (n-2)\widetilde{\theta}^\# \lrcorner d\beta + (n-1)\langle \beta, \widetilde{\theta} \rangle \widetilde{\theta} + (\widetilde{\delta} \beta) \widetilde{\theta} \right),$$

where all operators of the right hand side are associated to the metric \widetilde{g} . Applying this formula for $\beta = \widetilde{\alpha}$, the pull-back of α , and taking into account that $\widetilde{\alpha}$ is closed and co-closed and orthogonal on $\widetilde{\theta}$, we obtain that $\widetilde{\alpha}$ is also harmonic with respect to the Kähler metric, *i.e.* $\Delta^K \widetilde{\alpha} = 0$. Since on a Kähler manifold the Laplace operator commutes with the complex structure, it follows that $\Delta^K \widetilde{J}\widetilde{\alpha} = \widetilde{J}\Delta^K \widetilde{\alpha} = 0$. Applying now (14) to $\beta = \widetilde{J}\widetilde{\alpha}$ and projecting on M , we obtain:

$$(15) \quad \Delta J\alpha + (n-1)d(\langle J\alpha, \theta \rangle) + (n-2)\theta^\# \lrcorner dJ\alpha + (n-1)\langle J\alpha, \theta \rangle \theta + (\delta J\alpha)\theta = 0,$$

which further simplifies, using (13) and the fact that $\langle J\theta, \alpha \rangle = c$ is constant, to:

$$(16) \quad \Delta J\alpha + (n-1)c\theta + (\delta J\alpha)\theta = 0.$$

Taking in (16) the scalar product with θ and integrating over M yields $(n-1)c = 0$. Since $n \geq 2$, we get $0 = c = \langle J\theta, \alpha \rangle$, so $J\alpha$ is pointwise orthogonal to θ . Applying the codifferential to (16) yields

$$\delta d\delta J\alpha - \theta^\#(\delta J\alpha) = 0.$$

On the other hand, using the fact that $\theta^\#$ is Killing and holomorphic, together with (12), we obtain $\theta^\#(\delta J\alpha) = \mathcal{L}_{\theta^\#}\delta J\alpha = 0$. Hence, we get

$$\Delta \delta J\alpha = \delta d\delta J\alpha = 0.$$

Thus, the function $\delta J\alpha$ is constant, and as its integral over M vanishes, we get that $\delta J\alpha$ is identically zero. Substituting this into (16) implies that $J\alpha$ is harmonic.

(ii) Let $\beta \in \mathcal{H}^1(M)$. By Lemma 5.1, the function $\langle \beta, \theta \rangle$ is constant and thus $\beta - \langle \beta, \theta \rangle \theta$ is harmonic. According to (i), this completes the proof. \square

Remark 5.3. A direct consequence of Lemma 5.2 is the fact that the first Betti number of a compact Vaisman manifold is odd, which was proven in [12].

Proposition 5.4. *Let (M^{2n}, J, g) be a compact toric Vaisman manifold. Then the following assertions hold:*

- (i) *The lcK rank of $(J, [g])$ equals 1.*
- (ii) *M is obtained as the mapping torus of a compact toric Sasakian manifold.*

Proof. (i) The rank of any lcK structure on M is less than or equal to the first Betti number of M . Hence, it is enough to show that $H_1(M, \mathbb{R}) \cong \mathbb{R}$. According to the Hodge theory on compact manifolds and to the decomposition (11), it suffices to show that $\mathcal{H}_0^1(M) = \{0\}$.

Let $\beta \in \mathcal{H}_0^1(M)$ and let ξ be a fundamental vector field of the toric action. By Lemma 5.1 applied to β and to the Killing vector field ξ , the function $\beta(\xi)$ is constant. On the other hand,

ξ is twisted Hamiltonian, so there exists $h \in \mathcal{C}^\infty(M)$, such that $(J\xi)^{\flat} = \xi \lrcorner \Omega = d^\theta h = dh - h\theta$. We now integrate over M the constant $\beta(\xi)$:

$$\int_M \beta(\xi) dv = \int_M \langle J\beta, (J\xi)^{\flat} \rangle dv = \int_M \langle J\beta, dh - h\theta \rangle dv = \int_M (\delta(J\beta)h - h\langle J\beta, \theta \rangle) dv = 0,$$

where for the last equality we used that $J\beta \in \mathcal{H}_0^1(M)$ cf. Lemma 5.2, so $J\beta$ is orthogonal to θ and co-closed. Hence, $\beta(\xi) = 0$. In particular, the same holds also for $J\beta$, since $\mathcal{H}_0^1(M)$ is J -invariant, by Lemma 5.2.

Let us now consider a basis of the Lie algebra of the torus T^n , $\{\xi_1, \dots, \xi_n\}$. We denote by the same symbols the corresponding fundamental vector fields on M . According to the principal orbit theorem, there exists a dense open set M_0 of M , on which $\{\xi_1, \dots, \xi_n, J\xi_1, \dots, J\xi_n\}$ forms a basis of the tangent bundle of M_0 . As shown above, each $\beta \in \mathcal{H}_0^1(M)$ vanishes on $\{\xi_1, \dots, \xi_n, J\xi_1, \dots, J\xi_n\}$, hence $\beta = 0$ on M_0 , and by density also on M . We thus conclude that $\mathcal{H}_0^1(M) = \{0\}$.

(ii) We recall that the minimal covering \widehat{M} of a Vaisman manifold M , endowed with the Kähler metric g^K , is biholomorphic and isometric to the Kähler cone over a Sasakian manifold S . Moreover, each homothety of $(\mathbb{R} \times S, g^K)$ is of the form $(t, p) \mapsto (t + a, \psi_a(p))$, for some $a \in \mathbb{R}$ and ψ_a an isometry of S . If the Vaisman structure is assumed to be toric, then by [17, Theorem 4.9], the Sasakian manifold S is also toric.

Since by (i), the lcK rank of the toric Vaisman structure equals 1, the deck transformation group of the minimal covering \widehat{M} is $\Gamma \cong \mathbb{Z}$. Using this identification, we assume that Γ is generated by the strict homothety $(t, p) \mapsto (t + 1, \psi_1(p))$. Hence, M is obtained as the mapping torus $[0, 1] \times S / \sim$, where $(0, p) \sim (1, \psi_1(p))$. Moreover, this also shows that S is compact. \square

Remark 5.5. According to [16, Theorem 1.1], the fundamental group of a compact toric Sasakian manifold is a finite abelian group. Hence, from the proof of Proposition 5.4, (ii), it follows that the universal covering of a compact toric Vaisman manifold is the Kähler cone over a compact toric Sasakian manifold. We recall that compact toric contact manifolds were classified by E. Lerman in [15, Theorem 2.18].

6. THE KODAIRA DIMENSION OF TORIC LcK MANIFOLDS

Let (M, J) be a compact complex manifold of complex dimension n and $K := \Lambda^{n,0}M$ its canonical line bundle. We recall that the Kodaira dimension of (M, J) is defined as

$$\kappa(M) := \limsup_{\ell \rightarrow \infty} \frac{\log(\dim H^0(M, K^{\otimes \ell}))}{\log \ell},$$

where $H^0(M, K^{\otimes \ell})$ denotes the space of holomorphic sections of $K^{\otimes \ell}$. For properties of the Kodaira dimension, as well as for the Kodaira-Enriques classification of compact complex surfaces we refer to [2] or [13], [14].

Theorem 6.1. *The Kodaira dimension of a compact toric strict lcK manifold is $-\infty$.*

Proof. Let $(M^{2n}, J, [g])$ be a compact toric strict lcK manifold. An effective action of the torus T^n has n -dimensional principal orbits, and their union is a dense open set M_0 in M by the principal orbit theorem (see *e.g.* [6, Theorem 2.8.5]). Consequently, there exist real holomorphic vector fields X_1, \dots, X_n which mutually commute and such that X_1, \dots, X_n are linearly independent on M_0 . Since X_1, \dots, X_n are twisted Hamiltonian, Lemma 3.6 and Proposition 3.9 show that $\Omega(X_j, X_k) = 0$ for every $j, k \leq n$. Consequently $X_1, \dots, X_n, JX_1, \dots, JX_n$ are linearly independent on M_0 . Moreover, these $2n$ vector fields mutually commute, since X_1, \dots, X_n are holomorphic and mutually commute. We thus obtain holomorphic sections $Z_j := X_j - iJX_j$ of $T^{1,0}M$ satisfying $[Z_j, Z_k] = [Z_j, \bar{Z}_k] = 0$ for every $j, k \in \{1, \dots, n\}$ and such that Z_1, \dots, Z_n are linearly independent on M_0 . The anti-canonical bundle $(K_M)^* \simeq \Lambda^n(T^{1,0}M)$ thus has a non-trivial holomorphic section $\sigma := Z_1 \wedge \dots \wedge Z_n$.

Assume, for a contradiction, that some positive power $(K_M)^{\otimes k}$ of the canonical bundle has a non-trivial holomorphic section α . Then $\alpha(\sigma^{\otimes k})$ is a non-trivial holomorphic function on M , thus it is a non-zero constant. Consequently, σ is nowhere vanishing, so $\{Z_j\}_{1 \leq j \leq n}$ is a basis of $T^{1,0}M$ at each point of M . We denote by $\omega_j \in \Omega^{1,0}M$ the dual basis and claim that ω_j are closed holomorphic 1-forms. Indeed, since $\omega_i(Z_j) = \delta_{ij}$ and $\omega_i(\bar{Z}_j) = 0$, we obtain for all i, j, k :

$$\begin{aligned} d\omega_i(Z_j, Z_k) &= Z_j(\omega_i(Z_k)) - Z_k(\omega_i(Z_j)) - \omega_i([Z_j, Z_k]) = 0, \\ d\omega_i(Z_j, \bar{Z}_k) &= Z_j(\omega_i(\bar{Z}_k)) - \bar{Z}_k(\omega_i(Z_j)) - \omega_i([Z_j, \bar{Z}_k]) = 0. \end{aligned}$$

Define now the real $(1, 1)$ -form $\Omega := i \sum_{j=1}^n \omega_j \wedge \bar{\omega}_j$. From the above we have that Ω is closed. Moreover, the symmetric bilinear form $\Omega(\cdot, J\cdot)$ is positive definite, so Ω is a Kähler form on M compatible with J . By a result of I. Vaisman, [22, Thm. 2.1], the lcK structure is then globally conformally Kähler, contradicting the assumption that the lcK structure is strict.

This contradiction shows that the positive tensor powers of K_M have no holomorphic sections, hence $\kappa(M) = -\infty$. \square

7. TORIC LCK SURFACES

Using the Kodaira classification of compact complex non-Kählerian surfaces, Theorem 6.1 and a recent result of N. Istrati [10], we will now describe all compact complex surfaces carrying a toric strict lcK structure. Let us first recall the definition of Hopf surfaces.

Definition 7.1. A primary Hopf surface is a complex surface $(\mathbb{C}^2 \setminus \{0\})/\Gamma$, where Γ is the group generated by

$$\gamma(z_1, z_2) := (\beta z_1, \alpha z_2 + \lambda z_1^m),$$

with $\alpha, \beta, \lambda \in \mathbb{C}$, satisfying $0 < |\alpha|, |\beta| < 1$, $\lambda(\alpha - \beta^m) = 0$ and $m \in \mathbb{N} \setminus \{0\}$. The following two cases are distinguished:

- (i) The diagonal primary Hopf surfaces, when $\lambda = 0$.
- (ii) The non-diagonal primary Hopf surfaces, when $\lambda \neq 0$ (and thus $\alpha = \beta^m$).

A secondary Hopf surface is a finite quotient of a non-diagonal primary Hopf surface.

The main result of this section is the following:

Theorem 7.2. *The only compact complex surfaces admitting a toric strict lcK structure are the diagonal Hopf surfaces.*

Proof. Let (M, J) be a compact complex surface carrying a toric strict lcK structure. By Theorem 6.1, $\kappa(M) = -\infty$. The Kodaira classification of non-Kählerian compact complex surfaces shows that the only possible examples with Kodaira dimension $-\infty$ are the Inoue surfaces, the Hopf surfaces, the surfaces in the class VII_0^+ (i.e. with $b_2 > 0$), and blow-ups thereof (see e.g. [13] or [1, Proposition 1]). On the other hand, by a recent result of N. Istrati [10], any toric lcK manifold admits a toric Vaisman structure. By the classification of compact Vaisman complex surfaces, obtained by F. Belgun [1, Theorem 1], it follows that the only ones with Kodaira dimension $-\infty$ are the diagonal Hopf surfaces.

Let $M_{\alpha,\beta} := (\mathbb{C}^2 \setminus \{0\})/\Gamma$ be a diagonal Hopf surface, where Γ is the group generated by $\gamma(z_1, z_2) := (\beta z_1, \alpha z_2)$, according to Definition 7.1.

A Vaisman metric on $M_{\alpha,\beta}$ was constructed by P. Gauduchon and L. Ornea [8], as follows: let $\phi_{\alpha,\beta} : \mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{R}$ be the function implicitly defined as the unique solution of the equation

$$|z_1|^2 |\beta|^{-2\phi_{\alpha,\beta}(z)} + |z_2|^2 |\alpha|^{-2\phi_{\alpha,\beta}(z)} = 1,$$

for any $z \in \mathbb{C}^2 \setminus \{0\}$. It can be checked that the function $|\alpha\beta|^{\phi_{\alpha,\beta}}$ is a potential for a Kähler metric $g^K := dd^c |\alpha\beta|^{\phi_{\alpha,\beta}}$ on $\mathbb{C}^2 \setminus \{0\}$ and $g := |\alpha\beta|^{-\phi_{\alpha,\beta}} g^K$ projects to a Vaisman metric on $M_{\alpha,\beta}$ (for details see [8, Proposition 1]).

We now prove that the above defined Vaisman structure on $M_{\alpha,\beta}$ is toric. We first notice that $\mathbb{C}^2 \setminus \{0\}$ carries a T^2 -action given by

$$(17) \quad T^2 \longrightarrow \text{Diff}(\mathbb{C}^2 \setminus \{0\}), \quad (t_1, t_2) \cdot (z_1, z_2) = (t_1 z_1, t_2 z_2),$$

which is effective, holomorphic and Γ -invariant and hence descends to an effective, holomorphic T^2 -action on $M_{\alpha,\beta}$. The potential $\phi_{\alpha,\beta}$ is smooth and satisfies $\phi_{\alpha,\beta} \circ \gamma = \phi_{\alpha,\beta} + 1$, for any $\gamma \in \Gamma$, and is constant along the T^2 -orbits, so T^2 acts isometrically on $\mathbb{C}^2 \setminus \{0\}$ with respect to g^K . As the conformal factor between g and g^K is exactly this potential, it follows that T^2 also acts isometrically with respect to g and is a subgroup of $\text{Aut}_s(M_{\alpha,\beta})$. By Lemma 3.7, the T^2 -action is twisted Hamiltonian. \square

An alternative proof of Theorem 7.2 can be obtained without using the above cited result of N. Istrati, as a direct consequence of Theorem 6.1 and the following more general result, which is interesting in itself:

Proposition 7.3. *The only non-Kählerian compact complex surfaces of Kodaira dimension $-\infty$ admitting an effective holomorphic T^2 -action are Hopf surfaces.*

For the proof of Proposition 7.3, we will need the following standard result which relates holomorphic vector fields on a complex manifold and on its blow-up at some point (see for instance [4]):

Lemma 7.4. *Let M be a compact complex manifold. The holomorphic vector fields on the blow-up of M at a point p are exactly the lifts of holomorphic vector fields on M that vanish at p .*

Proof of Proposition 7.3. As mentioned above, by the Kodaira classification, the only non-Kählerian compact complex surfaces of Kodaira dimension $-\infty$ are the following: the Inoue surfaces, the Hopf surfaces, the surfaces in the class VII_0^+ (i.e. with $b_2 > 0$), and blow-ups thereof.

The space of holomorphic vector fields is at most one-dimensional on Inoue surfaces (see [9, Prop. 2 (ii), Prop. 3 (ii) and Prop. 5] or [1, Prop. 12]) and on surfaces in the class VII_0^+ (see e.g. [5, Remark 0.2]). Consequently, these surfaces cannot admit an effective holomorphic T^2 -action. By Lemma 7.4, the same holds for each of their blow-ups.

We now consider the two types of Hopf surfaces.

1) Diagonal Hopf surfaces and blow-ups thereof.

Let $M_{\alpha,\beta} := (\mathbb{C}^2 \setminus \{0\})/\Gamma$ be a diagonal Hopf surface, where Γ is the group generated by $\gamma(z_1, z_2) := (\beta z_1, \alpha z_2)$. Let us first note that $M_{\alpha,\beta}$ admits an effective holomorphic T^2 -action, according to (17).

In order to prove that a blow-up at a point of $M_{\alpha,\beta}$ does not admit an effective holomorphic T^2 -action, we start by determining the space of holomorphic vector fields on $M_{\alpha,\beta}$. Namely, we have the following

Claim. The space of Γ -invariant holomorphic vector fields on $\mathbb{C}^2 \setminus \{0\}$ is generated by

- a) $\left\{ z_1 \frac{\partial}{\partial z_1}, z_2 \frac{\partial}{\partial z_2}, z_2 \frac{\partial}{\partial z_1}, z_1 \frac{\partial}{\partial z_2} \right\}$, if $\alpha = \beta$.
- b) $\left\{ z_1 \frac{\partial}{\partial z_1}, z_2 \frac{\partial}{\partial z_2}, z_2^m \frac{\partial}{\partial z_1} \right\}$, if $\alpha = \beta^m$ for some integer $m \geq 2$.
- c) $\left\{ z_1 \frac{\partial}{\partial z_1}, z_2 \frac{\partial}{\partial z_2}, z_1^m \frac{\partial}{\partial z_2} \right\}$, if $\beta = \alpha^m$ for some integer $m \geq 2$.
- d) $\left\{ z_1 \frac{\partial}{\partial z_1}, z_2 \frac{\partial}{\partial z_2} \right\}$, if $\alpha \neq \beta^k$ and $\beta \neq \alpha^k$ for any $k \in \mathbb{N}$.

Proof of the Claim. Like before, any holomorphic vector field on $\mathbb{C}^2 \setminus \{0\}$ can be extended to \mathbb{C}^2 , since the singularity is isolated. Let $X := u \frac{\partial}{\partial z_1} + v \frac{\partial}{\partial z_2}$ be a holomorphic vector field, where u and v are two holomorphic functions on \mathbb{C}^2 . The vector field X is Γ -invariant if and only if $u(\alpha z_1, \beta z_2) = \alpha u(z_1, z_2)$ and $v(\alpha z_1, \beta z_2) = \beta v(z_1, z_2)$. Equivalently, $a_{jk}(\alpha - \alpha^j \beta^k) = 0$ and $b_{jk}(\beta - \alpha^j \beta^k) = 0$ for all $j, k \in \mathbb{N}$, where a_{jk} and b_{jk} are the coefficients of the power series of u and v respectively. Since $0 < |\alpha|, |\beta| < 1$, it turns out that in each of the four possible cases, X is a linear combination of the vector fields as stated in the claim. \checkmark

The next step is to show that there are no two linearly independent commuting holomorphic vector fields which vanish at the same point. Note that it suffices to prove this for the cases a) and b) in the above claim.

Case a) Assume that $\alpha = \beta$ and let $X, Y \in \text{span}\left\{ z_1 \frac{\partial}{\partial z_1}, z_2 \frac{\partial}{\partial z_2}, z_2 \frac{\partial}{\partial z_1}, z_1 \frac{\partial}{\partial z_2} \right\}$, such that there exists $w \in \mathbb{C}^2 \setminus \{0\}$ with $X_w = Y_w = 0$. Hence, there exist $a \in \mathbb{C}^2 \setminus \{0\}$ and $c_j \in \mathbb{C}$, such that $X_z = (a_1 z_1 + a_2 z_2) \left(c_1 \frac{\partial}{\partial z_1} + c_2 \frac{\partial}{\partial z_2} \right)$ and $Y_z = (a_1 z_1 + a_2 z_2) \left(c_3 \frac{\partial}{\partial z_1} + c_4 \frac{\partial}{\partial z_2} \right)$. Therefore,

the Lie bracket of X and Y is given by

$$\begin{aligned} [X, Y]_z &= (a_1 z_1 + a_2 z_2) \left((a_1 c_1 + a_2 c_2) \left(c_3 \frac{\partial}{\partial z_1} + c_4 \frac{\partial}{\partial z_2} \right) - (a_1 c_3 + a_2 c_4) \left(c_1 \frac{\partial}{\partial z_1} + c_2 \frac{\partial}{\partial z_2} \right) \right) \\ &= (a_1 z_1 + a_2 z_2) (c_2 c_3 - c_1 c_4) \left(a_2 \frac{\partial}{\partial z_1} - a_1 \frac{\partial}{\partial z_2} \right). \end{aligned}$$

Thus, if X and Y commute, then $c_2 c_3 = c_1 c_4$, hence X and Y are linearly dependent.

Case b) Assume that $\alpha = \beta^m$ for some integer $m \geq 2$ and let $X, Y \in \text{span}\{z_1 \frac{\partial}{\partial z_1}, z_2 \frac{\partial}{\partial z_2}, z_2^m \frac{\partial}{\partial z_1}\}$, such that there exists $w \in \mathbb{C}^2 \setminus \{0\}$ with $X_w = Y_w = 0$. This readily shows that there exist $a, b \in \mathbb{C}^2 \setminus \{0\}$ such that either

$$\begin{aligned} X_z &= a_1 z_2^m \frac{\partial}{\partial z_1} + a_2 z_2 \frac{\partial}{\partial z_2}, & Y_z &= b_1 z_2^m \frac{\partial}{\partial z_1} + b_2 z_2 \frac{\partial}{\partial z_2} \quad \text{or} \\ X_z &= (a_1 z_1 + a_2 z_2^m) \frac{\partial}{\partial z_1}, & Y_z &= (b_1 z_1 + b_2 z_2^m) \frac{\partial}{\partial z_1}. \end{aligned}$$

Therefore, we compute

$$[X, Y]_z = m(a_2 b_1 - a_1 b_2) z_2^m \frac{\partial}{\partial z_1}, \quad \text{respectively } [X, Y]_z = b_1 X_z - a_1 Y_z.$$

In both cases, X and Y commute if and only if they are linearly dependent.

By applying Lemma 7.4, we conclude that no blow-up of $M_{\alpha, \beta}$ carries two linearly independent commuting holomorphic vector fields, and in particular cannot carry an effective holomorphic T^2 -action.

2) Non-diagonal Hopf surfaces and blow-ups thereof.

Since the secondary Hopf surfaces are finite quotients of non-diagonal primary Hopf surfaces, it suffices to show that the latter ones do not admit an effective holomorphic T^2 -action. Let $M := (\mathbb{C}^2 \setminus \{0\})/\Gamma$ be a non-diagonal primary Hopf surface, where Γ is generated by $\gamma(z_1, z_2) := (\beta z_1, \beta^m z_2 + \lambda z_1^m)$, according to Definition 7.1. We first determine the space of holomorphic vector fields on M . These correspond to the holomorphic vector field on $\mathbb{C}^2 \setminus \{0\}$ invariant under the action of Γ .

Claim. The space of Γ -invariant holomorphic vector fields on $\mathbb{C}^2 \setminus \{0\}$ is generated by

$$X_1 := z_1 \frac{\partial}{\partial z_1} + m z_2 \frac{\partial}{\partial z_2}, \quad X_2 := z_1^m \frac{\partial}{\partial z_2}.$$

Proof of the Claim. Let $X = u \frac{\partial}{\partial z_1} + v \frac{\partial}{\partial z_2}$ be a Γ -invariant holomorphic vector field on $\mathbb{C}^2 \setminus \{0\}$, where u and v are two holomorphic functions on $\mathbb{C}^2 \setminus \{0\}$. Since the singularity is isolated, u and v extend holomorphically to the whole complex plane \mathbb{C}^2 . The Γ -invariance of X reads

$$(18) \quad u \circ \gamma(z) = \beta u(z),$$

$$(19) \quad v \circ \gamma(z) = m \lambda z_1^{m-1} u(z) + \beta^m v(z),$$

for all $z \in \mathbb{C}^2$. On the other hand, by induction on $n \in \mathbb{N}$, we establish that

$$(20) \quad \gamma^n(z) = (\beta^n z_1, \beta^{mn} z_2 + n \lambda \beta^{m(n-1)} z_1^m).$$

In the sequel, we use the fact that for every $\beta \in \mathbb{C}$ with $0 < |\beta| < 1$, $k \in \mathbb{N}$ and $n \in \mathbb{N} \setminus \{0\}$, an entire function f , which satisfies $f(\beta^n z) = \beta^k f(z)$ for any $z \in \mathbb{C}$, is either a monomial of degree $\frac{k}{n}$, if n divides k , or identically zero otherwise.

For $z_1 = 0$, (18) reads $u(0, \beta^m z_2) = \beta u(0, z_2)$. By the above remark, the function $z_2 \mapsto u(0, z_2)$ is linear if $m = 1$, or identically zero otherwise. Hence, there exists $c_m \in \mathbb{C}$ (which vanishes if $m \neq 1$), such that $z_1 = 0$ is a zero of the function $z_1 \mapsto u(z_1, z_2) - c_m z_2$. Thus $\tilde{u}(z) := \frac{u(z) - c_m z_2}{z_1}$ is an entire function and by (18) it satisfies $\tilde{u} \circ \gamma = \tilde{u} - \frac{c_m \lambda}{\beta}$. Therefore, $\tilde{u} \circ \gamma^n(z) = \tilde{u}(z) - n \frac{c_m \lambda}{\beta}$ for every $z \in \mathbb{C}^2 \setminus \{0\}$. Taking the limit in this identity for $n \rightarrow \infty$, and using the fact that $\gamma^n(z)$ tends to 0 for every z by (20), it follows that $c_m = 0$ and \tilde{u} is constant. We deduce that $u(z) = cz_1$ for some constant $c \in \mathbb{C}$. Substituting in (19) and iterating, we obtain by immediate induction

$$(21) \quad v \circ \gamma^n(z) = \beta^{nm} v(z) + nmc\lambda \beta^{m(n-1)} z_1^m, \quad \forall n \in \mathbb{N}.$$

In particular, for $n = 1$ and $z_1 = 0$, we readily obtain $v(0, \beta^m z_2) = \beta^m v(0, z_2)$, for all $z_2 \in \mathbb{C}$. So, there exists $c' \in \mathbb{C}$, such that $v(0, z_2) = c' z_2$, hence the entire function $z_1 \mapsto v(z_1, z_2) - c' z_2$ vanishes at $z_1 = 0$. Therefore, either $v(z) = c' z_2$, for any $z \in \mathbb{C}^2$, or there exists $k \in \mathbb{N} \setminus \{0\}$ and an entire function \tilde{v} , such that $v(z) - c' z_2 = z_1^k \tilde{v}(z)$ and $\tilde{v}(0, a) \neq 0$, for some $a \in \mathbb{C}$. In the first case, (19) implies that $c' = mc$. We now consider the second case. By (21), the function \tilde{v} satisfies

$$\tilde{v} \circ \gamma^n(z) = \beta^{(m-k)n} (\tilde{v} + n\lambda \beta^{-m} (mc - c') z_1^{m-k}).$$

We now let $n \rightarrow \infty$ and distinguish the following cases:

- if $k > m$, then $\tilde{v} \equiv 0$ and $c' = mc$.
- if $k < m$, then the function \tilde{v} vanishes at $(0, z_2)$, for all $z_2 \in \mathbb{C}$. This contradicts the definition of k and \tilde{v} .
- if $k = m$, then $c' = mc$ and \tilde{v} is constant.

We conclude that $u(z) = cz_1$ and $v(z) = mcz_2 + c'' z_1^m$, for some constants $c, c'' \in \mathbb{C}$. This proves the claim. \checkmark

A direct consequence of the above Claim is that there are no two linearly independent commuting Γ -invariant holomorphic vector fields on $\mathbb{C}^2 \setminus \{0\}$ which vanish at the same point. Hence, by Lemma 7.4, no blow-up of M carries an effective holomorphic T^2 -action.

Since the vector fields X_1 and X_2 commute, the flow of any holomorphic vector field $X := aX_1 + bX_2$, with $a, b \in \mathbb{C}$, is the composition of the flows of aX_1 and bX_2 , which are given by

$$\varphi_t(z_1, z_2) = (e^{at} z_1, e^{mat} z_2), \quad \psi_t(z_1, z_2) = (z_1, z_2 + btz_1^m).$$

Namely, the flow of X is $(\varphi_t \circ \psi_t)(z_1, z_2) = (e^{at} z_1, e^{mat}(z_2 + btz_1^m))$. Hence, the orbits of X are all relatively compact in $\mathbb{C}^2 \setminus \{0\}$ if and only if $a \in i\mathbb{R}$ and $b = 0$, *i.e.* X is a multiple of X_1 . On the other hand, on $M := (\mathbb{C}^2 \setminus \{0\})/\Gamma$, the projection of the vector field $Y := \log \beta \cdot X_1 + \frac{\lambda}{\beta^m} \cdot X_2$, where \log denotes a branch of the complex logarithm containing β in its domain of definition, is a holomorphic vector field with closed orbits. Consequently,

the real parts of the projections of X_1 and Y to M induce an effective holomorphic T^2 -action on M . This finishes the proof Proposition 7.3. \square

Remark 7.5. Let us note that Theorem 7.2 can be deduced from Proposition 7.3 as follows. Let (M, J) be a compact complex surface admitting a toric lcK metric g . By Theorem 6.1 and Proposition 7.3, (M, J) is a Hopf surface. It thus remains to rule out the case of non-diagonal Hopf surfaces. We assume now that $M := (\mathbb{C}^2 \setminus \{0\})/\Gamma$ is a non-diagonal primary Hopf surface. Then, Lemma 3.6 implies that $T^2 \subseteq \text{Aut}_s(M, J, [g])$. Furthermore, by Proposition 4.4, the T^2 -action lifts to $\mathbb{C}^2 \setminus \{0\}$, which is not only the universal covering of M , but also the minimal covering of any lcK structure on M , since $\pi_1(M) = \mathbb{Z}$. This contradicts the last computation in the proof of Proposition 7.3, which shows that there do not exist two linearly independent holomorphic Γ -invariant vector fields on $\mathbb{C}^2 \setminus \{0\}$ with relatively compact orbits. We conclude that a non-diagonal primary Hopf surface does not carry a toric lcK structure. On the other hand, each diagonal Hopf surface admits a toric lcK structure, where the T^2 -action is defined by (17).

Note that any non-diagonal primary Hopf surface (M, J) admits an lcK metric (cf. [1, Proposition 11]), which can be averaged to an lcK metric g by an argument of L. Ornea and M. Verbitsky, [18], such that $T^2 \subseteq \text{Aut}(M, J, [g])$. However, as seen above, T^2 is never a subgroup of $\text{Aut}_s(M, J, [g])$.

REFERENCES

- [1] F. A. Belgun, *On the metric structure of non-Kähler complex surfaces*, Math. Ann. **317** (2000), no. 1, 1–40.
- [2] W. Barth, C. Peters, A. Van de Ven, *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Band **4**, Springer Verlag, Berlin, 1984.
- [3] A. L. Besse, *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008.
- [4] J. Carrell, A. Howard, C. Kosniowski, *Holomorphic vector fields on complex surfaces*, Math. Ann. **204** (1973), 73–81.
- [5] G. Dloussky, K. Oeljeklaus, M. Toma, *Surfaces de la classe VII admettant un champ de vecteurs, II*, Comment. Math. Helv. **76** (2001), 640–664.
- [6] J.J. Duistermaat, J.A.C. Kolk, *Lie Groups*, Universitext, Springer Verlag, Berlin, 2000.
- [7] P. Gauduchon, *La 1-forme de torsion d'une variété hermitienne compacte*, Math. Ann. **267** (1984), 495–518.
- [8] P. Gauduchon, L. Ornea, *Locally conformally Kähler metrics on Hopf surfaces*, Ann. Inst. Fourier **48** (1998), 1107–1127.
- [9] M. Inoue, *On Surfaces of Class VII₀*, Invent. Math. **24** (1974), 269–310.
- [10] N. Istrati, *A characterisation of toric LCK Manifolds*, arXiv:1612.03832.
- [11] T. Kashiwada, *On V-harmonic forms in compact locally conformal Kähler manifolds with the parallel Lee form*, Kodai Math. J. **3** (1980), 70–82.
- [12] T. Kashiwada, S. Sato, *On harmonic forms in compact locally conformal Kähler manifolds with the parallel Lee form*, Ann. Fac. Sci. Univ. Nat. Zaïre (Kinshasa) **6** (1980), no. 1-2, 17–29.
- [13] K. Kodaira, *On compact complex analytic spaces I*, Ann. Math. **71** (1960), 111–152; II, ibid. **77** (1963), 563–626; III, ibid. **78** (1963), 1–40.
- [14] K. Kodaira, *On the structure of compact complex analytic spaces I*, Am. J. Math. **86** (1964), 751–798; II, ibid. **88** (1966), 682–721; III, ibid. **90** (1969), 55–83; IV, ibid. **90** (1969), 1048–1066.
- [15] E. Lerman, *Contact toric manifolds*, J. Symplectic Geom. **1** (2003), 785–828.

- [16] E. Lerman, *Homotopy groups of K -contact toric manifolds*, Trans. of the Amer. Math. Soc. **356** (2004), no. 10, 4075–4083.
- [17] M. Pilca, *Toric Vaisman manifolds*, J. Geom. Phys. **107** (2016), 149–161.
- [18] L. Ornea, M. Verbitsky, *Automorphisms of Locally Conformally Kähler Manifolds*, Int. Math. Res. Not. **2012** (2012), no. 4, 894–903.
- [19] L. Ornea, M. Verbitsky, *LCK Rank of locally conformally Kähler manifolds with potential*, J. Geom. Phys. **107** (2016), 92–98.
- [20] K. Tsukada, *Holomorphic forms and holomorphic vector fields on compact generalized Hopf manifolds*, Compos. Math. **24** (1994), no. 1, 1–22.
- [21] I. Vaisman, *On locally conformal almost Kähler manifolds*, Israel J. Math. **24** (1976), 338–351.
- [22] I. Vaisman, *On locally and globally conformal Kähler manifolds*, Trans. Amer. Math. Soc. **262** (1980), no. 2, 533–542.
- [23] I. Vaisman, *Generalized Hopf manifolds*, Geom. Dedicata **13** (1982), 231–255.

FARID MADANI, INSTITUT FÜR MATHEMATIK, GOETHE UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STR. 10, 60325 FRANKFURT AM MAIN, GERMANY

E-mail address: madani@math.uni-frankfurt.de

ANDREI MOROIANU, LABORATOIRE DE MATHÉMATIQUES DE VERSAILLES, UVSQ, CNRS, UNIVERSITÉ PARIS-SACLAY, 78035 VERSAILLES, FRANCE

E-mail address: andrei.moroianu@math.cnrs.fr

MIHAELA PILCA, FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, UNIVERSITÄTSSTR. 31 D-93040 REGENSBURG, GERMANY *and* INSTITUTE OF MATHEMATICS “SIMION STOILOW” OF THE ROMANIAN ACADEMY, 21, CALEA GRIVITEI STR. 010702-BUCHAREST, ROMANIA

E-mail address: mihaela.pilca@mathematik.uni-regensburg.de